


$$\begin{array}{c} 1B \\ & \begin{array}{c} (2\text{-Ac}) \\ \text{Abe}\alpha 1 \\ & \uparrow \\ & 3 \\ \text{Man}\alpha 1 \rightarrow 4Rha\alpha 1 \rightarrow 3Gal}\alpha 1 \end{array} \\ & \begin{array}{c} (2\text{-Ac}) \\ \text{Abe}\alpha 1 \\ & \uparrow \\ & 3 \\ \rightarrow 2Man}\alpha 1 \rightarrow 4Rha\alpha 1 \rightarrow 3Gal}\beta 1 \rightarrow 4 = R \end{array}$$

an O-antigen or repeating polysaccharide



Heptaacyl Lipid A





FIGURE 2

3A Haemophilus influenzae strain 2019 LOS

Fig. 4 Western Blot Analysis of Anti LOS-Conjugate Sera Against LOS From Different Bacteria

Lane #



BRL lows Sample blk <i>N. mening</i> A1	NTHi TN106 NTHi P860295	N. gc 52407 P+ N. gc 1756 P+	H. pylori 1103H. pylori 43579	H. pylori 1105 M. cat 1230:359	M. cat 25238 M. cat 430:345	<i>N. gc</i> LB2 <i>N. gc</i> Pgh3-1
3 2 -	4 ፕ	9	ထတ	10 11	13	4 15